

Trawl fishery (southern inshore region) harvest strategy: 2021–2026



Business area owner Management & Reform

Endorsed by Deputy Director-General (Fisheries & Forestry) in accordance with delegated

powers under Part 2, Division 1 (Harvest Strategies) of the Fisheries Act 1994

Approved by Minister responsible for fisheries in accordance with section 16 of the Fisheries Act

1994

Revision history

Version no.	Approval date	Comments
0.01	September 2020	Draft harvest strategy for consultation
1.00	June 2021	Approved harvest strategy

Enquiries and feedback regarding this document can be made as follows:

Email: info@daf.qld.gov.au

Telephone: 13 25 23 (Queensland callers only)

(07) 3404 6999 (outside Queensland)

Monday, Tuesday, Wednesday and Friday: 8 am to 5 pm, Thursday: 9 am to 5 pm
Department of Agriculture and Fisheries GPO Box 46 BRISBANE QLD 4001 AUSTRALIA

Website: daf.qld.gov.au

Interpreter statement



Post:

The Queensland Government is committed to providing accessible services to Queenslanders from all culturally and linguistically diverse backgrounds. If you need an interpreter to help you understand this document, call 13 25 23 or visit daf.qld.gov.au and search for 'interpreter'.

© State of Queensland, 2021

The Queensland Government supports and encourages the dissemination and exchange of its information. The copyright in this publication is licensed under a Creative Commons Attribution 4.0 International (CC BY 4.0) licence.

Under this licence you are free, without having to seek our permission, to use this publication in accordance with the licence terms.



You must keep intact the copyright notice and attribute the State of Queensland as the source of the publication.

Note: Some content in this publication may have different licence terms as indicated.

For more information on this licence, visit https://creativecommons.org/licenses/by/4.0/.

The information contained herein is subject to change without notice. The Queensland Government shall not be liable for technical or other errors or omissions contained herein. The reader/user accepts all risks and responsibility for losses, damages, costs and other consequences resulting directly or indirectly from using this information.

Contents

What the harvest strategy is trying to achieve	4
Fishery overview	4
Stocks covered by the harvest strategy	5
Management units for the harvest strategy	5
Summary of management information	6
Fishery objectives	7
Catch shares	7
Managing the performance of the fishery	8
Management of target species	11
Management of secondary and by-product species	12
Management of ecological risks from fishing	13
Monitoring social and economic performance	14
Data collection, validation and assessment	15
Information and research priorities	15
Schedule of performance monitoring, assessment and review	16
Acronyms and definitions	18

What the harvest strategy is trying to achieve


This harvest strategy has been developed in line with the *Queensland harvest strategy policy* to manage trawl fishery resources within the southern inshore trawl region. Current stock levels for saucer scallops, the key species in the region, are overfished and rebuilding is required to achieve maximum sustainable yield (MSY). This harvest strategy aims to initially rebuild the biomass to MSY, with a long-term goal of achieving the maximum economic yield (MEY) of the resource. While saucer scallops are the target species that drives fishing effort in the southern inshore trawl region, other species including Moreton Bay bugs, banana prawns and tiger prawns are also harvested.

The aim of this harvest strategy is to manage fishing mortality through setting sustainable effort limits at a level that allows the stock to achieve its biomass targets. For all other retained species, effort triggers have been designed to monitor changes in fishing behaviour or stock trends, and hence optimising economic yield, while at the same time being precautionary in detecting changes in species composition within historic catch levels. Other management tools (e.g. size limits, spawning closures etc.) may also be used to support the sustainable management of stocks under this harvest strategy.

Fishery overview

In Queensland, the annual catch of saucer scallops has been declining since 2001. From 2014 to 2017, catch was less than 546 tonnes meat weight (2730 tonnes whole weight) per year, near the lowest MSY estimate across a range of productivity scenarios for the stock. A stock assessment estimated that spawning biomass of the east coast biological stock in 2015 may have been as low as 5–6% of the 1977 unfished level. Results of the 2017 fishery-independent survey of abundance also showed relatively low densities of pre-recruits and scallops older than one year.

At 134 tonnes annual meat weight (670 tonnes whole weight) in 2017, landings of saucer scallops by the east coast otter trawl fishery were at an historical low. This is likely due in part to recent management intervention, including total closure of high abundance scallop replenishment areas from November 2016, and prohibition on harvesting during the May—October spawning season. However, the annual catch rate in 2017 also decreased to the lowest level seen since 1997, when recruitment failed. Average monthly catch rates in late 2017 were about 40% higher than the historic lows of January 2015 to April 2016, providing some evidence of stock rebuilding at least in the southern-most part of the fishery.



A long-term decline in the annual number of scallop harvesting days has been evident since 1997, when the stock was first considered to be overfished and effort in 2017 was at an historically low level. However, a shift in fleet composition towards more efficient vessels has increased fishing power since 2000.

Stocks covered by the harvest strategy

While saucer scallops (which are subject to stock rebuilding) are the primary target species, this harvest strategy also manages a number of other permitted species that can be retained while operating under a T1 licence. Table 1 provides a summary of fish stocks covered by this harvest strategy.

Table 1: Summary of fish stocks covered by this harvest strategy

Feature	Details	
Target species (tier 1)	Saucer scallops (Amusium balloti)	
Secondary species (tier 2)	Brown and grooved tiger prawns (<i>Penaeus esculentus</i> and <i>Penaeus semisulcatus</i>) Banana prawns (<i>Fenneropenaeus merguiensis</i>) Moreton Bay bugs (<i>Thenus</i> spp.)	
Byproduct species (tier 3) Permitted trawl species: Balmain bugs blue swimmer crabs cuttlefish mantis shrimps octopus pipefish red champagne lobsters slipper lobsters threadfin bream three-spotted crabs		
Biology	Saucer scallops live to a maximum of four years old and 140 mm shell height 50% maturity is reached at 85–90mm shell height (~12 months)	

Management units for the harvest strategy

The single management unit for this harvest strategy is from latitude 22 degrees south down to the southern end of Hervey Bay, excluding the eastern king prawn fishing grounds. The fishery area is defined in Schedule 8 of the Fisheries (Commercial Fisheries) Regulation 2019.

Ballot's saucer scallop (*Ylistrum balloti*, formerly classified as *Amusium balloti*) is distributed from Israelite Bay in Western Australia, across the tropics, to the southern coast of New South Wales. There are two adjacent management units for saucer scallop stock – one from Queensland that is managed by Fisheries Queensland (latitude 22°–27° south) and another along the New South Wales coast that is managed by the New South Wales Department of Primary Industries. This harvest strategy manages the Queensland part of the scallop stock.

Summary of management information

A summary of the management arrangements for the southern inshore trawl region is set out in Table 2 (overleaf). Fishers may access copies of fisheries legislation at legislation.qld.gov.au or visit fisheries.qld.gov.au for the latest information on fishing rules.

Table 2: Summary of management arrangements for the southern inshore trawl region

Feature	Details	
Commercial access	Primary commercial fishing licence with a T1 fishery symbol	
Relevant fisheries legislation	Fisheries Act 1994	
	Fisheries (General) Regulation 2019	
	Fisheries (Commercial Fisheries) Regulation 2019	
	Fisheries Declaration 2019	
	Fisheries Quota Declaration 2019	
Other relevant legislation	Great Barrier Reef Marine Park Act 1975 and Great Barrier Reef Marine Park Regulations 2019 (Cwlth)	
	Marine Parks Act 2004	
	Environment Protection and Biodiversity Conservation Act 1999 and Environment Protection and Biodiversity Conservation Regulations 2000 (Cwlth)	
Harvest strategy workshop	Southern inshore region harvest strategy workshops are held at least annually	
	Further advice on proposed management arrangement and fishery performance will be shared with the trawl fishery working group Terms of reference and communiques are available at fisheries.qld.gov.au	
Gear	Otter trawl apparatus may be used, generally with quad gear configuration	
	Refer to <u>fisheries legislation</u> for specific gear requirements and rules	
Main management methods	Primary management method is individual effort units and a total allowable effort cap for the region Scallop-specific effort cap (Other management methods include: Ilimited access through primary commercial fishing licences 20 m maximum vessel length hull unit (HU) limit of 120 HU gear restrictions such as net length and mesh size	
	 no-take closure period for scallops (1 May – 30 November) spatial and temporal closures 	
Fishing year	20 November – 23 December, 3 January – 19 November	

Feature	Details
Stock status	Stock status is assessed using the nationally agreed <u>Status of Australian</u> <u>Fish Stocks</u> (SAFS) classification framework – saucer scallops are listed as 'depleted' (SAFS 2018)
	*Note: The classification system used as part of the SAFS reporting is assessed against a 20% biomass sustainability criteria. Therefore, although a species may be classified as 'sustainable' in SAFS, this does not mean that the biomass is meeting the targets set out in the <i>Queensland Sustainable Fisheries Strategy: 2017–2027</i> . For more species specific biomass estimates, consult the relevant stock assessment for that species.
Accreditation under the Environment Protection and Biodiversity Conservation Act 1999	Part 13: Accredited (expires 2021) Part 13A: Accredited (expires 2021) Visit environment.gov.au

Fishery objectives

The objective of the harvest strategy is to manage the fishery in accordance with the objectives of the *Fisheries Act 1994* and the *Queensland Sustainable Fisheries Strategy: 2017–2027*.

Fishery objectives set out the aspirations and operational direction for the management of this fishery. The primary objective of the southern inshore trawl region fishery is to:

 maintain the target species in the southern inshore trawl region at, or returned to, a target spawning biomass level of MSY for the fishery.

In pursuing the primary objective, the harvest strategy aims to:

- minimise and mitigate any unacceptable ecological risks arising from fishing-related activities
- maximise economic performance of the commercial sector
- monitor the broader social and economic benefits of the fishery to the community.

Catch shares

This harvest strategy aims to maintain the existing catch shares between sectors. The resource allocation arrangements set out in Table 3 ensure that catch shares among sectors are maintained in response to changes in the total allowable commercial effort (TACE). The existing resource allocation arrangements (as at 2018) are set out in Table 3 and this harvest strategy will aim to maintain the existing catch shares between the sectors.

The traditional fishing rights of Aboriginal peoples and Torres Strait Islanders are protected under native title legislation and relate to harvest for domestic, communal and non-commercial purposes. Accordingly, traditional and customary fishing is recognised in Queensland and is not a defined allocation.

Aboriginal peoples and Torres Strait Islanders and their communities continue to express a desire to have more economic opportunities through fishing, particularly in their own sea country. The *Aboriginal and Torres Strait Islander commercial fishing development policy* provides for an Indigenous fishing permit to be issued, on a case-by-case basis and in accordance with section 54 of the Fisheries (General) Regulation 2019, to provide opportunities to take part in fishing-related business.

Table 3: Resource allocation arrangements for the southern inshore trawl region

Species	Commercial fishing*	Recreational fishing** (including charter)	
Saucer scallops	99%	1%	

^{*} Commercial catch data is based on the existing commercial catch level.

Managing the performance of the fishery

Key indicators measure the fishery's performance. The indicators relate to the objectives and use reference points to establish acceptable performance (Table 4 overleaf). The indicators measure the relative amount of fish biomass of key stock(s) against target and other reference points. The default biomass reference points identified in this harvest strategy are:

- a target reference point (Btarg) of 40% of the unfished spawning biomass (for rebuilding the target species, scallops, to maximum sustainable yield (MSY)) being the relative biomass level the harvest strategy aims to achieve.
- a limit reference point of 20% of the unfished spawning biomass (Blim) being the biomass level that the harvest strategy aims to avoid If the stock is assessed to be below Blim, the risk to the stock is unacceptably high and the stock is defined as 'depleted'.

For key stocks, performance indicators and sustainable harvests for all sectors will be estimated from a stock assessment. The aim is to measure the capability for the stock to attain the target biomass level (Btarg 60%), and at which point the harvest strategy will be considered as meeting its fishery objectives.

The decision rules for setting a sustainable harvest in the southern inshore trawl region are based on a 'hockey stick' approach. This is where the TACE is set based on a linear relationship between Blim, where the level of fishing mortality (F) is equal to zero, and Btarg, where the exploitation rate and TACE are set at the level to achieve MEY (Figure 1).

The decision rule takes into account the current biomass level of the stock for determining the TACE to achieve the Btarg. The recommended TACE is calculated by applying the rate of fishing mortality to achieve Btarg to the current spawning biomass level. As a result, the recommended TACE represents the total catch from all sectors (including discards) that can be harvested in the next three years, to move the current biomass level towards the target level. A discount factor may also be included to account for uncertainty and to reduce the risk of a fishery not achieving its objectives.

If the spawning biomass of a stock falls below Blim, targeted fishing of the stock will cease and a rebuilding strategy be developed to rebuild the spawning biomass above Blim within a biologically reasonable timeframe (e.g. based on mean generation time¹) and informed by the Queensland Harvest Strategy Policy..

^{**} Recreational catch share includes charter fishing and is based on information from statewide recreational fishing surveys.

¹ a generation is defined as the average age of full maturity for the fish species.




Figure 1: Showing the 'hockey stick' rule – Blim is limit reference point, Bmey is the biomass at MEY, B0 is the unfished biomass at 100%, F is fishing mortality and Ftarg is the level of fishing mortality for Bmey

Setting total allowable commercial effort

In response to the spawning biomass estimate for scallops being below the limit reference point of 20%, a range of management changes have been proposed to further influence stock recovery in order to achieve the objectives set out for the scallop stock:

- extend existing no-take scallop closure in the southern inshore and southern offshore trawl regions from 1 May until 20 November
- implement southern inshore regional fishery closures from 20 September to 20 November and from 23 December to 3 January
- set a regional effort cap for scallops in the southern inshore trawl region at 80 000 effort units.

With no MSY estimate available, the current effort level of 246 000 effort units was used to set the initial effort cap commencing in 2020. In addition to the regional effort cap, a scallop-specific effort cap was set at recommended effort levels of 80 000 effort units.

The harvest strategy is also designed to adjust the TACE based on updated regional fishing power estimates. Fishing power will be updated periodically as part of each scheduled stock assessment. When an updated stock assessment becomes available that indicates the average level of fishing power has changed, then the TACE will be set at a level to achieve Btarg.

Management of secondary commercial and by-product species

If biomass is not available as a primary indicator for secondary (tier 2) and by-product (tier 3) species, commercial catch will be monitored to assess changes in fishing mortality. Annual commercial catch triggers are used to assess changes in fishing mortality when compared to historic catch levels.

Secondary species (tier 2) are important species that aren't always targeted by fishers. These species have historical commercial catch ranges set and annual commercial catch is monitored against these. Historical commercial catch ranges from 2010 to 2019 were used to set the indicator values (Table 4). As the level of exploitation changes outside of historic levels, species will be elevated to higher levels of monitoring, assessment and management.

By-product species (tier 3) are sometimes retained by fishers and have trip limits or no-catch limits in place. Annual commercial catch of by-product species will be monitored against a two-year mean of commercial catch to detect changes in fishery behaviour that may represent unacceptable risk to sustainability. Mean commercial catch from 2017 to 2019 was used to set the reference point for by-product species.

Table 4: Performance indicators and reference points for the southern inshore trawl region

Species	Performance indicator	Reference point / buffer	Reference level
Scallops (tier 1 species)	Spawning biomass	Target (Btarg)	40% spawning biomass
Scallops	Spawning biomass	Limit reference point (Blim)	20% spawning biomass
Secondary and by-product species (if available) (tier 2 and tier 3)	Biomass	Target (Btarg)	60% biomass
Secondary and by-product species (if available) (tier 2 and tier 3)	Biomass	Limit reference point (Blim)	20% biomass
Secondary species (tier 2)	Annual commercial logbook catch	Commercial catch range	2010–2019
By-productsSpecies (tier 3)	Annual commercial logbook catch	2-year mean commercial catch	2017–2019

Management of target species

1.0 Decision rules for saucer scallops

The decision rules provide guidance to the TACE based on estimates of biomass being available. The decision rules use the outputs of the stock assessment and aim to rebuild the target stock to achieve a target biomass (Btarg) of 40%.

- 1.1 If the biomass is at or above Btarg, set the TACE at a level that maintains biomass at Btarg.
- 1.2 If biomass is below Btarg and above Blim, the TACE should be set as inferred by the hockey stick approach, where fishing mortality is reduced to the rate that allows the biomass to increase effectively back to Btarg.
- 1.3 If biomass is at or below Blim, there will be no targeted fishing for that species, and a rebuilding strategy will be developed to increase the stock biomass to above Blim within a biologically reasonable timeframe and as informed by the Queensland Harvest Strategy Policy..
- 1.4 If any new information becomes available indicating that the assessment and TACE-setting arrangements are not consistent with the sustainable management of the fishery, decision rules must be reviewed and, if appropriate, the reference points or timeframes should be adjusted.

Notwithstanding that:

1.5 The rate of fishing mortality should not exceed the level of fishing mortality required to maintain a stock at MSY at equilibrium.

2.0 Breakout rules for the take of scallops

- 2.1 If new information becomes available to suggest that a change to the fishery closure dates should be made to support rebuilding of the stock and meet the objectives of the fishery, then the fishery closure dates should be adjusted for the following season.
- 2.2 If and when any new information becomes available indicating that the assessment and TACE-setting arrangements are not consistent with the sustainable management of the fishery, the scientific method and review rules must be reviewed and, if appropriate, the reference points must be adjusted.

3.0 Decision rules for fishing power adjustments

To ensure that the TACE reflects current fishing power in the southern inshore trawl region, the decision rules allow for adjustment to the TACE if a new fishing power estimate becomes available. The new fishing power estimate will be calculated as the mean change in the most recent five years of fishing power estimates.

- 3.1 If no new estimate of fishing power is available, then the existing estimate is applied to the TACE. If the fishing power estimate is to be applied at regular intervals between scheduled stock assessment years, the TACE will be set to adjust accordingly.
- 3.2 When a new estimate of fishing power becomes available, generally every three years through the scheduled stock assessment, then the TACE will be adjusted to the new estimate of fishing power (i.e. an increase in fishing power will result in a proportional decrease in the TACE).

Management of secondary and by-product species

4.0 Decision rules for secondary species

The secondary species are classified as tier 2 species in this fishery and do not have catch limits. The harvest control rules below monitor effort shift to ensure there is no unacceptable levels of fishing pressure for tier 2 species (e.g. tiger prawns and Moreton Bay bugs).

The harvest strategy also includes decision rules to allow management arrangements to be implemented if updated biomass estimates become available.

- 4.1 If the annual harvest is between 3 and 11 tonnes for tiger prawns, or 130 and 350 tonnes for Moreton Bay bugs, then no management action is required.
- 4.2 If the harvest for two consecutive years is outside of the catch range of 3–11 tonnes for tiger prawns, or 130–350 tonnes for Moreton Bay bugs, a review is to be undertaken to understand the reason for the increased harvest, assess the risks and ensure catch of a species does not increase more than 20% above the upper catch range. If rule 4.2 is triggered. management action must be in place for the following fishing season until a detailed review is completed (e.g. trip limits, size limits or spatial/temporal closures). If the review identifies that a species is of increasing importance, the species may be considered for further stock assessment, monitoring or management action. If the review identifies sustainability is at risk, a stock assessment for this species is required within three years. If a stock assessment becomes available, then the stock assessment will be used to inform management of the species

5.0 Decision rules for by-product species

By-product species are classified as tier 3 species in this fishery and do not have catch limits. The harvest control rules below monitor effort shift to ensure there is no unacceptable levels of fishing pressure for tier 3 species (e.g. Balmain bugs, blue swimmer crabs).

- 5.1 If the three-year average harvest of any species is no more than 10% above the average catch from 2017 to 2019, then no management action is required.
- 5.2 If the three-year average commercial catch of any species is more than 10% above average catch from 2017 to 2019, a review is to be undertaken to understand the reason for the increased harvest and assess the risks and ensure catch of a species does not exceed 10% above average catch from 2017 to 2019. If catch of a species exceeds more than 10% above average catch from 2017 to 2019, management action must be in place for the following fishing season until a detailed review is completed (e.g. trip limits, size limits or spatial/temporal closures). If the review identifies sustainability is at risk, a stock assessment this species is required within three years

6.0 Breakout rules for secondary and by-product species

6.1 If a biomass estimate is available through a stock assessment for secondary or by-product species that indicates a reduction in fishing mortality is required to achieve a Btarg or avoid Blim, then management action is to be taken (e.g. trip limits, size limits or spatial/temporal closures) to pursue the fishery objectives.

Management of ecological risks from fishing

A foundation of sustainable fisheries management is managing the impact of fishing activities on non-target species and the broader marine ecosystem. Ecological risk assessments (ERAs) identify and measure the ecological risks of fishing activity and identify issues that must be further managed under harvest strategies. The decision rules below are in place to minimise and mitigate high ecological risks arising from fishing.

- 7.1 If an ERA identifies fishing impacts that are considered to generate an unacceptable level of risk to any ecological component, a review is triggered to investigate the reason for the risk , and appropriate management action taken to reduce the risk to an acceptable level.
- 7.2 If the southern inshore trawl region effort footprint in any given year is greater than the 2019 effort footprint, a review will be undertaken to understand the reason for the increased effort footprint and identify appropriate management strategies to reduce the risk, including options that reduce the permitted area that can be trawled. If the review identifies sustainability is at risk, management action is required within two years.

The most recent ERA was completed in 2012 through a collaborative project between the Great Barrier Reef Marine Park Authority, Fisheries Queensland and the Queensland Seafood Industry Association. The ERA was for the entire Great Barrier Reef Marine Park area; however, the key findings for the southern inshore trawl region are as follows:

- Risks from trawling have reduced in the Great Barrier Reef since the introduction of a fishery management plan in 1999 and subsequent management actions.
- Marine park zoning has been important in reducing risks from trawling.
- Commercial fishers have been proactive in seeking and using better fishing practices to reduce trawling impacts.
- Current risk levels from trawling activities are generally low. Under current practices and based on 2009
 effort levels, the overall ecological risks from trawling to harvested species, as well as the broader
 environmental values and integrity, are low.
- Trawl fishing effort is a key driver of ecological risk. Real risks will be addressed through harvest strategies and the fishery working group, taking into account the factors contributing to high-risk ratings, the adequacy of current management regime and, if applicable, strategies to reduce and mitigate the risk to the environment. Measures may include regional trawl effort caps under harvest strategies, further reductions in trawl bycatch, high compliance with rules, and accurate information from ongoing risk monitoring etc.

For more information about the ERA for the east coast otter trawl fishery in the Great Barrier Reef Marine Park, refer to the <u>summary report</u> or the <u>full technical report</u>.

Fisheries Queensland developed the <u>Ecological risk assessment guideline</u> to assess ecosystem impacts of fishing activities. Future risk assessments will be undertaken in line with the guideline to reassess any current or new ecological risks that may arise in the fishery. ERAs can be undertaken more frequently if there are significant changes identified in fishery operations, management activities or controls that are likely to result in a change to previously assessed risk levels.

Monitoring social and economic performance

The *Queensland Sustainable Fisheries Strategy: 2017–2027* outlines the target to set sustainable catch limits based on achieving B_{MEY} (around 60% of unfished biomass) to support the most economically efficient use of the resource, improve the fishing experience for all sectors and promote a resilient system that can respond to other adverse environmental conditions (e.g. floods, cyclones and bleaching). The harvest strategy decision rules have been set up to achieve this target biomass level.

The objectives and performance indicators in Table 5 will be used to monitor the social and economic performance of this fishery. The management options outlined are intended to provide some guidance on options that could reasonably be considered alongside the decision rules if fishery trends are of concern.

Table 5: Social and economic indicators for the southern inshore trawl fishery

Objective	Performance indicators	Management options
Maximise economic performance of the commercial sector	 Potential indicators to monitor include: capacity utilisation catch per unit effort (average per day) costs, earnings and net financial and economic profit net economic returns, gross state production, gross value of production sale and lease price profit decomposition (using profit or lease price) to determine impacts of prices, costs and stock/catch rates on changes in profits 	Consider regulatory and non- regulatory options Adjust management as needed Options include minimum holding, latent effort review
Monitor the broader social and economic benefits of the fishery to the community	 Potential indicators to monitor include: fisher satisfaction (with their fishing experience – commercial and recreational) Recreational fisher participation and economic information percentage of unit/licences that are owned (rather than leased) Gini coefficient of unit owner (measure of concentration) percentage of total costs/inputs purchased from local businesses/residents income generated (crew plus profit – gross value added) proportion of catch sold locally fish prices number of platforms / number of active licences / total capacity community satisfaction (with their fisheries and the way in which they are managed) 	Consider regulatory and non-regulatory options Adjust management as needed
Maintain United States turtle exclusion device inspection program and Section 609 United States export accreditation	United States inspection report	Amend management and fisheries legislation as required to align gear controls with accreditation requirements

Data collection, validation and assessment

Fishery-dependent data (self-reported)

Catch and effort data is obtained through commercial logbook returns and real-time landing reports. The catch and effort data required to determine the standardised commercial catch rate for key species is obtained from catch and effort logbook returns and vessel tracking data. Commercial catch rates are standardised to account for fishing power, along with a range of potential influencing variables. The trawl logbook is available at business.qld.gov.au.

Fishers are also required to report any interactions with protected species in a mandatory threatened, endangered and protected animal logbook.

Fishery-dependent data (independent validation)

All commercial fishing vessels are required to have vessel tracking systems installed and active on their vessels. Vessel tracking data is used to verify effort information reported in commercial fishing logbooks. As an effort-managed fishery, compulsory effort unit deductions provide an accurate record of fishing activity.. Queensland Boating and Fisheries Patrol undertake routine and intelligence-based at-sea and landing (unload) inspections to check compliance and validate reported information.

Fishery-independent information

Fisheries Queensland conducts an annual fishery-independent trawl survey for saucer scallops. The survey samples 0+ and 1+ scallop age classes, which can be used to calculate a relative abundance index.

Scientific assessment of stock

Since 2017, a collaborative investment by the Queensland Government, Fisheries Research and Development Corporation and universities has been delivered annually to support monitoring and assessment of scallop stocks and to inform rebuilding strategies. An updated stock assessment was completed in 2019 (including data up until 2018) and provided MSY and MEY catch and effort estimates for scallops.

Information and research priorities

Key information and research priorities have been identified in Table 6 to help meet the objectives of this harvest strategy. These will be reviewed and updated as required through the harvest strategy workshop.

Table 6: Information and research priorities for the southern inshore trawl region

Project description	Explanation of need	Priority
Regular periodical fishing power surveys	Collect information to improve catch rate standardisation	Medium
Bycatch reduction device (BRD) testing and evaluation program to support continued innovation	To support continued innovation of trawl BRDs	High

Schedule of performance monitoring, assessment and review

Annual performance monitoring and assessment

The fishery's performance will be monitored against this harvest strategy **annually**. This will include an annual harvest strategy workshop to provide operational advice on the fishery's performance and any matters that may need addressing.

The primary performance measure is spawning biomass, which will be used to review the TACE every three years. In the intervening years, fishing power estimates will be applied to the TACE, thus reducing the allowable effort proportionally. Refer to the most current <u>stock assessment</u>.

While harvest strategies provide certainty and transparency in terms of management decisions in response to certain fishery information, there must also be flexibility to allow new information or changing circumstances to be appropriately considered. There may be instances in which a stock assessment may need to be available prior to, or delayed beyond, the scheduled date. Any change to the stock assessment schedule should be considered by the harvest strategy workshop and decided on by the chief executive based on the below conditions:

- If during the period between scheduled stock assessments the chief executive is concerned that a performance indicator (e.g. stock status, standardised commercial catch rate, total harvest) suggests the stock is not performing in a way that will achieve the target biomass level, the chief executive may decide that a stock assessment will be undertaken before the scheduled timeframe.
- If the chief executive is satisfied that (1) indicators for the stock suggest it is achieving, or rebuilding to, target biomass levels, and that there is a low ecological risk to the stock under the current management arrangements, or (2) if resourcing requirements prohibit the ability for an assessment to be delivered in the scheduled timeframe, the chief executive may decide that a scheduled stock assessment will be delayed.

Table 7: Schedule of performance monitoring, assessment and review

	Year 1	Year 2	Year 3	Year 4	Year 5
Assessment program	Modelled stock assessment	Catch and effort monitoring	Catch and effort monitoring	Modelled stock assessment	
Management program	Review TACE, reference points and fishing rules Fishing power adjustment is required	Review of catch and effort data Adjust TACE for fishing power Bring forward management decisions if needed	Review of catch and effort data Adjust TACE for fishing power Bring forward management decisions if needed	Review TACE, reference points and fishing rules Fishing power adjustment is required	Harvest strategy review

Harvest strategy review

This harvest strategy will remain in place for a period of five years, after which time it will be fully reviewed in accordance with the *Fisheries Act 1994*. The harvest strategy may also be subject to further review and amendment as appropriate within the five-year period if any of the following circumstances arise:

- there is new information that substantially changes the status of a fishery, leading to improved estimates of indicators relative to reference points
- drivers external to management of the fishery increase the risk to fish stock/s
- it is clear the strategy is not working effectively and the intent of the harvest strategy policy is not being met.

For more information on the processes for amending harvest strategies, refer to the <u>Queensland harvest</u> <u>strategy policy</u>.

Acronyms and definitions

Acronym/term	Definition		
Biomass	Total weight or volume of a stock or component of a stock (e.g. spawning stock biomass would refer to all adult (reproductively mature) fish in a population)		
Biomass limit reference point (B <i>lim</i>)	The point beyond which the risk to the stock is regarded as unacceptably high		
Biomass target (Btarg)	The desired biomass of the stock		
BRD Bycatch reduction device			
Bycatch	 A species that is incidentally either: taken in a fishery and returned to the sea killed or injured as a result of interacting with fishing equipment in the fishery, but not taken Bycatch can include protected species 		
By-product	Any part of the catch that is kept or sold, but is not the target species By-product makes some contribution to the value of the catch in a fishery but less than that of key commercial species		
Catch-per-unit-effort	The number or weight of fish caught by a unit of fishing effort Can be used as an index of relative abundance or indicator of change in the fishery		
Ecological risk assessment (ERA)	An assessment process that evaluates the relative risk posed by fishing on species, habitats and communities within a fishery		
F	Fishing mortality		
Ftarg	Fishing mortality target		
Maximum economic yield (MEY)	Sustainable level of harvest that allows net economic returns (profit) to be maximised		
Maximum sustainable yield (MSY)	The maximum average sustainable annual fishing mortality that can occur on a stock over an indefinite period under prevailing environmental conditions		
SAFS	Status of Australian Fish Stocks		
Total allowable commercial catch (TACC)	The harvest limit set for the commercial fishing sector usually achieved through setting TACC, but sometimes through input controls		
Total allowable commercial effort (TACE)	The annual effort limit set for a stock, species or species group, used to control commercial fishing mortality within a fishery		